
**A DETAILED ANALYSIS ON SUBSTANCE
MISUSE AMONG DIFFERENT ETHNICITIES
IN THE USA**

by Maisha Rafa



United International University
QUEST FOR EXCELLENCE

PROJECT

**A Detailed Analysis on Substance Misuse
Among Different Ethnicities in the USA.**

Submitted By:

Maisha Rafa

ID: 111182090

Major: MIS

School of Business & Economics

United International University

Submitted To:

Mr. Abdullah Al Fahad

Lecturer

School of Business &
Economics

United International
University

Date of Submission: 07-06-24

Letter of Transmittal

7th June, 2024

Mr. Abdullah Al Fahad
Lecturer
School of Business and Economics
United International University

Dear sir,

Subject: **Submission of Thesis Title: “A Detailed Analysis of Substance Misuse Among Different Ethnicities in the USA”**

I am pleased to submit my thesis titled “A Detailed Analysis of Substance Misuse Among Different Ethnicities in the USA” for obtaining a Bachelor of Business Administration (BBA) degree at United International University.

The research presented in this thesis investigates examining the patterns and trends of substance misuse among ethnic populations of the USA to identify the disparities and root causes for suggesting successful intervention strategies.

I believe that the findings of this research contribute valuable insights to the field of my research.

I appreciate very much your support and guidance throughout this research.

I kindly submit this thesis for evaluation. I look forward to any feedback and advice you might give me.

Thank you very much.

Yours sincerely,

Maisha Rafa
111182090
Management of Information Systems (MIS)
School of Business & Economics
United International University

Acknowledgment

At the very beginning, I would like to thank Almighty Allah that I have been able to complete my thesis paper successfully.

I would also like to thank my distinguished faculty, Mr. Abdullah Al Fahad, for his exceptional guidance, support, and encouragement, through which he was able to inspire me. A thesis like this would be almost impossible without his professional advice and invaluable comments.

My friends, who inspired me all the way, without their moral support and feedback, this would not have been possible. Special thanks to UIU for giving me the resources to complete this thesis.

And lastly, I cannot ignore the extensive support and understanding of my family. The patience, love, and belief in me to finish this academic pattern were the challenges that ran me through.

With all my heart, I worked to deliver my best effort by fulfilling all requirements of my respective faculty and gaining insights about the following thesis. This educational thesis may serve as a lesson to get knowledge, give insights, and present structured analysis to the readers on the subject.

Declaration

I, Maisha Rafa, do hereby declare that the thesis provided is a result of my very own research and ideas. In the course of this academic exercise, I owe so much to everyone who, in one way or another, gave me immense support. Any contribution towards my study is gratefully acknowledged and referenced. This work has not been previously presented for any other degree. I confirm that all ethical considerations and academic standards were highly observed in the entire research process.

Maisha Rafa
111182090
Management of Information Systems (MIS)
School of Business & Economics
United International University

Abstract

This deep look into the substance abuse of different ethnic groups in the US sees the complex web of variables that determine the influence of drug use patterns within different groups. Based on an enormous pool of empirical data and research studies, this extensive study threads through the intricate interactions between urban neighborhood settings, violence exposures, socioeconomic differences, cultural norms, and systemic factors that create patterns of substance abuse seen in different ethnic groups. Based on sophisticated research using both quantitative and qualitative methods, this research provides an in-depth and comprehensive understanding of the base mechanisms leading to differences in substance abuse in the American setting.

This will shine a light on the broader consequences of substance abuse affecting different communities beyond illustrating the drug use pattern differences. The outcomes of the analysis shall be useful to researchers, legislators, and other interested parties who need to devise holistic interventions dealing with the root causes of substance abuse for the promotion of health outcomes at a societal level. This paper will unlock the door for more comprehensive strategies that aspire to diminish substance misuse issues and lead to a healthier, more resilient society for all people by being cognizant of the complex web of influences that shape drug use behaviors and the divergent needs of the different ethnic groups.

Keywords:

Drug, Substance Abuse, Drug Misuse in the USA

Contents

Acknowledgment.....	ii
Declaration	iii
Abstract	iv
1. Introduction:	1
2. Literature Review	2
3. Methodology.....	17
4. Analysis.....	19
5. Findings	20
6. Limitations.....	23
7. Recommendations	24
8. Discussion.....	26
9. Conclusion.....	27
References	28

1. Introduction:

Substance misuse in the United States is a complex issue which has various levels of impact on different ethnicities and diverse communities. Through many former researches, Black communities have been found to be more involved in substance misuse and are more affected by it than Whites. (Beth A. Reboussin, 2018). In later researches, it has also been found that along with race, socioeconomic status plays a huge role in substance consumption. In another older research the patterns of substance misuse has been classified as follows: Non-Hispanic Whites, who are less prone to sanctions, were found to be more into drugs and alcohol consumptions; on the other hand, such practices were found to be the least among Asian students; and lastly, Asian-White multiracial people were found to be at the midpoint between Asians and White. (Martin, 2019). There is a complex interaction of social determinants such as discrimination, socioeconomic status, and cultural norms that necessitates tailored interventions for the underlying drivers of substance misuse disparities among ethnic backgrounds. Former research has identified that substance misuse can also be impacted by adverse childhood experiences and peer influences. (Stritzel, 2021). Youth from minority communities who have lived around toxic or violent social environments are more likely to continue the trajectory of Marijuana. (Beth A. Reboussin, *The Impact of the Urban Neighborhood Environment on Marijuana Trajectories During Emerging Adulthood*, 2018). Furthermore, the studies found there are gender differences in the patterns of substance misuse, with men indicating more substance use as opposed to women, especially in college facilities.

In this context, a significant part of former research has focused on the role of adverse childhood experiences (ACEs) in the eventual development of substance abuse behaviors. According to existing results, it was found that adverse experiences, like psychological abuse and family substance abuse, are more prevalent for girls, while boys are more likely to face physical abuse. Some risk factors varied in statistical differences among boys and girls. (Lindsay Lebana, 2020). For girls, lower self-control showed higher risk of substance misuse while depression among boys was an indicator of higher substance misuse. (Lindsay Lebana, 2020). Among both boys and girls, criminality among peers showed higher chances of substance abuse. (Lindsay Lebana, 2020). Neighborhood has also proved to play a significant part in substance use/abuse. Substance misuse was also found to be intricately correlated to concerns of poverty, including homelessness, unemployment and also incarceration. (Karen A. D'Angelo, 2019). In light of these, there is a need to understand that the multifaceted influences on substance misuse, including age, gender, socioeconomic status, and environmental factors, have a pivotal role in developing targeted interventions and support systems that will address the complex challenges associated with drug abuse among diverse ethnic populations in the United States.

This study takes a more in-depth look at the various dynamics within drug use behaviors in its attempt to unravel the complex interplay of factors that lead to substance abuse among diverse ethnic communities. Based on the perspectives of previous researchers, the current study explores the disparities, barriers, and determinants that affect the rates of substance misuse

within minority populations, emphasizing the increased influence of drug abuse within foreign communities.

2. Literature Review

2.1 (Beth A. Reboussin, 2018)

In this research, the authors looked into the influence of neighborhood environment in the formation of drug misuse trajectories among young adults in urban settings. The number of alcohol outlets in the neighborhood, crime rate, and level of poverty are strong predictors of substance use among young adults. The neighborhood factors exert very important influences in the trajectory of shaping substance use during the period of emerging adulthood. Theoretically, studies such as Garberino's socially toxic environment posit that unsafe neighborhood conditions spur on high-risk behaviors, one of them being drug use.

Neighborhood environments significantly influence the pathways to drug use among young adults in urban settings. Addressing the issues of social disadvantage, neighborhood characteristics, and social role transitions intrinsic in developing effective interventions is key to reducing disparities in substance abuse across ethnicities in the USA. The study makes an effort to analyze the complex interplay of factors in order that researchers and policymakers would be in a position to work towards promoting healthier outcomes among young adults in urban neighborhoods.

2.2 (Nisha Behariera, 2018)

The authors have conducted a detailed study on the relationship between exposure to violence and substance abuse behaviors over time. Data from National Longitudinal Study of Adolescent to Adult Health in Add Health program (a nationally representative dataset) was used to examine the behaviors among adolescence and adults. The data collected sheds light upon how exposure to violence can impact substance use in long term. The study measured correlations between measures of exposure to three forms of violence in adolescence and substance use outcomes assessed in both adolescence and adulthood. The research also measured individual associations with the substances of cocaine and methamphetamines.

The results of the study suggested strong associations between exposure to violence through adolescence and substance misuse. The findings conclude that experiencing direct violence showed strong impact on substance abuse in adulthood, while witnessing violence (indirect exposure) had stronger impact in adolescence. Both experiencing and witnessing severe violence alarms the rise of Post-Traumatic Stress Disorder diagnosis. Maladaptive practices like substance abuse have resulted to be found as a coping mechanism for such trauma victims

as the substances may provide temporary relief from distress or numb painful emotions associated with traumatic events.

2.3 (Harvey L. Nicholson, 2018)

In their specified demographic research of prescription opioid misuse authors have focused on differences in prescription opioid misuse among Black adults in the US. The study showcases a framework of substance abuse patterns among a specific demographic in the US. This research uses a quantitative research methodology, which uses the data from National Survey on Drug Use and Health (NSDUH). The authors used weighted logistic regression models arranged by gender and race. Total samples from 60,133 respondents were accumulated to compare data of POM in both Black men and women were compared to that of both White men and women.

The results revealed that Black women with a high school degree had significantly lower odds of POM compared to having received less. Living in relatively rural areas also showed a lower odd of POM. In addition, being aged between 35 and 49 compared to being aged between 18 and 25 decreased odds of POM. On the other hand, participation in one or more government programs increased odds of POM. Both White and Black women who had a history of substance misues were also found to be at a higher risk of POM. The overall results stated that in comparison to Black women, White women were not more likely to engage in POM. However, it was also noted that White men were more likely to engage in POM compared to Black men.

2.4 (Monnat, 2018)

In his paper, a rural sociology and public health expert has researched whether drug mortality rates in different locations were mostly influenced by opioid supply or socioeconomic distress.

The findings of the study showed that both the opioid supply factors and socioeconomic distress were significant drivers of mortality due to substance misuse from rural to urban locations. The highest non-Hispanic white drug mortality rates were in large metro areas and most rural regions were found to have fewer deaths. All the data may vary due to different socioeconomic conditions like poverty, familial distress etc. but the results of the study showed that the drug mortality rates were significantly influenced by opioid supply.

2.5 (Veronica A. Peara, 2018)

This paper discussed the role of social factors of prescription opioid deaths and heroin overdose deaths across diverse states. Hierarchical Bayesian space time models were used in the study. Four main models were weighed, two of those evaluated POD and the other two evaluated HOD. Likelihood ratio tests were performed to assess whether the added interaction terms improved fit. Economic characteristics that were assessed included poverty, percentage of

unemployed adults, percentage of residents with a high school degree or less and percentage of Non-Hispanic White males.

Moreover, the association between the four economic characteristics and POD did not differ across levels of urbanicity, however, the relationship between economic stressors and HOD differed quite significantly at different levels of urbanicity. In more urban regions, HOD was localized in communities typified by poverty, unemployment, and/or low median household income. In contrast, only lower educational attainment was associated with higher HOD in the most rural counties. While economic stressors are related to POD in both rural and urban areas, they appear to assume a more influential role in HOD within urban areas. Results of the study depicted that social determinants for overdose of heroin may differ across the rural and urban areas.

2.6 (Martin, 2019)

The author has conducted a research examining the relationship between socioeconomic status and drug and alcohol consumption for the affluent collegian United States. The study found that individuals with high SES were at a higher risk of alcohol and drug consumption. He used self-report data based on a large sample of undergraduates with multi-level modeling applied for logistic and ordered logistics models in the analysis, followed by making weighted analysis.

Positive correlations were observed between parental educational attainment and SES and substance use. Whites were discovered to have the highest prevalence of substance abuse, while Asians had the lowest prevalence. It was determined that Asian-White biracial students were intermediate between Asians and Whites. Compared to students who resided in a fraternity or sorority, those who resided with their parents were less likely to use substances. A reduced rate of substance consumption was found among married individuals.

2.7 (Karen A. D'Angelo, 2019)

Substance misuse is a very complex public health issue. The photovoice study was conducted to understand the perspective of individuals who belonged to the communities that have been disproportionately affected. It reflects community views on the association of place and health in relation to substance misuse in Hartford, Connecticut. Photovoice was applied in three stages: the focus groups, training and photography, and the last stage was photo analysis and discussion. The participants were trained in the protocols of the study, how to use the camera and safety. Participants were also provided with digital cameras, alongside information sheets pertaining to the study and photo consent forms. This paper utilizes a combination of member checking and thematic analysis of the photographs.

In this study, three major themes included perception of the participants of environmental risk factors, strategies to cope in such a challenging environment and recommendations from the participants for prevention and treatment of substance abuse. The participants characterized the city as "drug-infested" because of the wide promotion and availability of both legal and

illicit drugs. It was also concluded that substance abuse was linked with poverty, homelessness, unemployment and incarceration. The use of illegal drugs and sale of those in public places were described as the major reinforcement factors of substance misuse. Substance abuse was found to be a strong health-related concern.

2.8 (Lindsay Lezana, 2020)

The research looked at how substance use and criminal misconduct during adolescence among males and females are associated with adverse childhood experiences. The methods that would be used to carry out such a study would include data analysis from the PHDCN; a prospective longitudinal design with three waves of data collection, using multiple imputations for missing data and applying two-level HGLM to estimate the fixed effects of ACEs on substance use and criminality.

Boys' delinquency and girls' substance use were found to be uniquely associated with adverse childhood experiences, according to the results of the Hierarchical Generalized Linear Models. The relationship between adverse childhood experiences and subsequent delinquency did vary for boys and girls: the impact of adverse childhood experiences on boys' delinquency was maintained as significant even after statistical controls while the impact on girls' delinquency became nonsignificant with statistical controls. Likewise, the impact of adverse childhood experiences on substance use differed for boys and girls in that exposure to adverse childhood experiences increased girls' chances of substance use, while with the other risk factors controlled, its impact on boys' substance use turned out to be nonsignificant.

2.9 (Elizabeth D. Nesoff, 2019)

The researchers have conducted a study to compare the geographic dispersal of fentanyl-involved and non-fentanyl-involved fatal overdoses from 2014 to 2018 in Cook County, Illinois. It provides insights on the distinguishing geographic distribution of deaths that involved fentanyl overdoses and confirms the speculation that states that the drug markets in resource-deprived areas rely severely on sale of fentanyl. This study's methodology involves a spatial analysis of the locations of overdose fatalities, which includes fentanyl and non-fentanyl opioids and poly-drug use. The analysis also includes a logistic regression analysis and a mapping of the locations of overdose. For analysis, the authors relied on data from the Cook County Medical Examiner's Office dating from 2014 through 2018. The data contain detailed toxicology reports and GPS coordinates, showing the exact location of overdose throughout Cook County, Illinois.

Non-fentanyl overdoses were less likely to cluster in geographic space than fentanyl overdoses, which presented a significantly higher risk in certain communities. Male sex, Black and Latino race/ethnicity, and younger age were identified as substantially increasing the likelihood of a fentanyl-related overdose. The only neighborhood predictor was found to be the neighborhood deprivation.

2.10 (Tamara Rushovich, 2020)

The authors have conducted a study to characterize the local opioid epidemic in comparison to the national epidemic, describe the association between neighborhood economic hardship and race/Hispanic ethnicity and the rate of opioid-related overdose death in Chicago, and delineate demographic trends and the communities with the highest rates of opioid-related overdose death. Methodology of this research was using data from the American Community Survey. The analysis of various demographic variables was used, and the Chicago rates compared to national rates, mapping out opioid-related overdose deaths.

These results have concluded that Chicago has a higher rate of overdose deaths related to opioids compared to national findings, with a different distribution by demographics, particularly in neighborhoods facing higher rates of poverty. Death rates due to opioid overdose among NH Black Chicagoans were higher than those of NH White and Hispanic Chicagoans. The correlation between opioid-related overdose mortality and economic hardship varied by racial/Hispanic ethnicity subgroup. This suggests that structural causes of economic hardship should be considered when addressing the opioid crisis.

2.11 (Gregory Phillips II, 2020)

In this research, the researchers have recorded higher substance uses among sexual and gender minorities compared to heterosexual and cisgender individuals. The research design used for the current study was a cohort study in which 488 young FAB SGM people took part, with an accelerated longitudinal design combining two cohorts: a late-adolescent and a young adult sample. The predictor variables were Engagement with LGBTQ-specific community, Current gender identity, Sexual orientation, and Degree of openness about being non-heterosexual to people around the participants. The dependent variables were uses of alcohol, marijuana and non-medical prescription drug and Alcohol-associated problems. The measured variables were age, ethnicity, gender identity, sexual orientation, openness about sexual minority identity, Places visited to be around other LGBTQ young people, History with LGBT-specific victimization, Non-prescription drug use, Cigarette usage, Use of prescription drugs, Alcohol use disorders. The authors tested if substance misuse and victimization rates differed in those who shared spaces with LGBTQ communities than those who did not.

The study found noteworthy correlations between substance misuse and victimization. LGBTQ spaces were also found to be associated with both adaptive and maladaptive coping functions. People who shared spaces with LGBTQ communities were found to be more exposed to victimization as well.

2.12 (Leslie D. Williams P. U., 2020)

This study estimates HIV prevalence among people who inject drugs in 89 of the US metropolitan areas that were highly populated. Mixed effects models with time nested within metropolitan statistical areas were used to estimate HIV prevalence among people who inject drugs. In their calculation of annual estimates of HIV prevalence, they used literature review estimates, regressed over a number of data series, and computed MSA-specific annual estimates of HIV prevalence between 1992 and 2013. The confirmation of the analysis was done through National HIV Behavioral Surveillance data. Analyses were computed by combining data from multiple sources and adjusting the model using baseline estimates. The researchers also examined the trends in the patterns by geographic region, number of persons tested for HIV, and baseline HIV prevalence for showing time trends and variability of trends in HIV prevalence among PWID.

Decreases in HIV prevalence among people who inject drugs was found through 2002. It was followed by a period of stabilization and steep increases after 2010. An average increase in HIV prevalence was found in the later years. Contradictorily, prevalences in the Eastern US were found to decrease in all years.

2.13 (Jacqueline Korpicsa, 2020)

It is a study that provides a summary of trends in substance use, sexual health, violence and victimization, and suicide risk among Chicago Public High School (CPS) students. This research mined data from the YRB survey by the CDC and Prevention among high school students in Chicago. The four domains were categorized as substance use, violence and victimization, sexual health risk and suicide risk. Findings were adjusted using the Stata program.

Significant improvement is reflected in health risk exposure among public high school students in Chicago during the 20-year period. Use of substances including cigarettes and other illegal drugs were found to have reduced.

2.14 (Jasmine Drake, 2020)

Authors have identified the opioid epidemic cost in the White, Black, and Hispanic communities in the United States, but more particularly, on the government and culturally sensitive policy considerations regarding racial disparities in response. Deaths were analyzed by year, state, substance type, and race and ethnicity. Deaths from opioid substance overdose were selected based on ICD10 underlying cause-of-death codes, and data was pulled from CDC WONDER Online Multiple Cause of Death database by the authors. Age-adjusted rates were used in an attempt to make comparisons between groups with different age distributions more equitable.

According to this study, the cases of deaths due to opioid overdose suddenly surged for Blacks and Hispanics. The Blacks, non-Hispanics, rose from 1130 in the year 1999 to 5513 in the year 2017, and Hispanics rose from 1058 in 1999 to 3932 in 2017. There were approximately 399,230 cases of opioid-related deaths nationally from 1999 to 2017 in the United States. The White, Non-Hispanic population accounted for around 323,939 deaths, compared to 75,291 by all other races.

2.15 (Jerel M. Ezell, 2020)

The article discusses the rural opioid crisis, systemic factors and social forces that impact drug use, rising rates of drug injection, the stigma associated with it and its implications on mental and physical health, and the isolation of individuals with stigmatized markers. Taken together, the approaches of this research article include semi-structured interviews, audio-recording of the interviews, analyzing data inductively, guaranteeing reliability by coding of two coders, and approval by the Institutional Review Boards and informed consent by all participants. The dataset used in this study is the Delta Rural Health Study.

The results indicated that opioid use outside medical practice in rural areas was associated with low-employment opportunities and high levels of physically demanding jobs, which predisposed individuals to chronic pain and opioid prescription. Other results were that stakeholders had mixed views of opioid use and people who use drugs, which were mostly placed within the framework of morality and character, and labeling of people who used drugs along perceived social and illness traits. The results also pointed out stigmatizing at the micro-level as the use of colorful labels and stereotypes.

2.16 (Leslie D. Williams M. E.-A., 2021)

The present paper addresses the interaction of two common social factors: stigma associated with substance-use and injection-related social norms as predictors for hepatitis C infection. The baseline data were selected for an ongoing longitudinal study being conducted with young injectors from Chicago suburbs and members of their injection risk network. To examine the relationships between HCV infection, safer injection norms, and substance use-related stigma, an adjusted logistic regression model was used.

Results of the study showed HCV infection playing a critical role in the effect of the stigma related to substance use on safer injection norms. Studies find a strong correlation between HCV infection and the interaction of safer injection norms with risks for HCV and substance-use-related stigma.

2.17 (Marika Sigal, 2021)

In this study, the researchers have given a dive into the association in relationship between parental monitoring and adolescent substance use in Latino and Black adolescents. This study states that neighborhood risks and peer influences on minor and major substance use provide a framework for contextual risk factors to understand the substance use in different ethnic groups.

In this study, the methodology followed is collecting data from adolescents through self-report surveys in North Carolina, doing path analyses and multigroup models, getting approval from an institutional review board, collecting data from three types of schools, and ensuring accuracy through trained research assistants and data checks.

The results showed maternal monitoring to be directly related to reduced exposure to substance misuse and criminal activities. Both maternal and paternal monitoring in Latinos and Blacks were directly related to gateway substance use. This analysis found that substance use was indirectly related to exposure to neighborhood crime and drugs by way of peer victimization and acquaintance delinquency.

2.18 (Lin Liu, 2021)

Authors of this study have conducted this study to show the how deteriorated family bonding and economic hassles may influence individuals to resume drug use after incarceration. Looking into the effects of private and public levels of social control on post-release drug use from this study could provide insights on how challenges of reentry after being released from imprisonment could lead to a relapse of substance abuse. The methodology used in the study included a longitudinal study conducted by the Urban Institute to explore the process of prisoner reentry.

Within the last year of being released, 22% had used illicit drugs, or one in five. Neighborhood cohesion ranked considerably higher within communities offering employment, medical coverage, and financial aid programs. It was further discovered that family exerted a significant influence in inhibiting illicit drug use. However, marriage, which has been cited by many as a predictor that reduces crime, did not serve to protect the respondents from illicit drug use. Combined, the personal, parochial, and public domains of social control were in place to insulate the released prisoners from the use of illicit drugs. Community cohesion was also found to inhibit released prisoners' drug use.

2.19 (Hanlin Zhou, 2021)

This paper discusses the association between places that involve drugs and factors related to place management and accessibility. It used Google Street View imagery to assess the environment and provides insights into the impact of safety score, traffic lights, poles, traffic signs, and roads on drug activities in Cincinnati, Ohio. The insights on the drug places given in this paper may contribute to broader research into drug abuse. The methodology involves the use of Google Street View imagery to calculate an aggregate safety score, extraction of physical elements at the address location, use of the binary logistic regression model, standardization of the independent variables, and checking for multicollinearity. The researchers compared the street view environment of the drug-related incidents to that of the street robbery incidents.

The results found that safety score, traffic lights, and poles had significantly negative influence on drug activities, while traffic signs and roads have shown statistically significant positive impacts. It was also found that the buildings had positive impacts. The drug locations were therefore strongly related to a safety score and various streetscape variables.

2.20 (Cara Jane Bergo, 2021)

In this study, the researchers have provided a context of the opioid crisis and its association with HIV outbreaks that are related to injection drug use. The methodology used includes a multivariable negative binomial regression and collection of ZIP code level indicators from various sources.

The opioid epidemic and injection drug use contributed to outbreaks of HIV in communities nationwide. High-ranking ZIP codes of vulnerability were dispersed throughout the state, but concentrated in the rural southern region of the state. Sub-county-level measurement of risk enabled more targeted surveillance and prevention efforts in high-risk communities.

2.21 (Anna Hotton, 2021)

It made use of random-effects meta-regression as a method for trend analysis in homelessness and injection risk behaviors over time, abstracting de-identified data from eleven research studies, data harmonization, and analysis. Data on persons who inject drugs came from the CDPH NHBS PWID cycles 2014, CIDUS 2 &, local studies administered by UICCOI investigators.

Quite to the contrary, there was a dramatic trend for increasing homelessness among young IDUs over time, mirroring the trend for increasing juvenile homelessness in the overall population. No overall changes in hazardous behavior over time, although homelessness among YP-IDUs was associated with injection risk behaviors and exchange sex.

2.22 (Jeffery Sauer a, 2021)

This study outlines the opioid overdose epidemic in light of social determinants of health in relation to getting healthcare services. The methods used in this study included global and local Moran's I analysis; nonspatial and spatial models; and spatiotemporal Bayesian hierarchical Poisson models.

The results detected a strong positive spatial autocorrelation in ED visits for heroin, methadone, and cocaine in 2019. The maps showed quite distinct spatial disease risk surfaces for each substance in the greater Baltimore metropolitan statistical area.

2.23 (Stritzel, 2021)

There have been many previous studies that have proved adverse childhood experiences to be linked to future substance misuse. Current research, which has evoked debate, addresses the degree to which peer and community contexts influence adolescent substance misuse. Therefore, it sheds light on the ways through which social and environmental aspects determine drug use. The methodologies used derive from negative binomial regression and path analysis using Mplus and the multilevel random intercept model. The design consists of two major elements: the Community Survey, which is administered to acquire neighborhood cluster-level data, and the Longitudinal Cohort Study, whose format was divided into three phases to acquire individual data.

More than half of the sample reported verbal and physical abuse. Many of the youth reported the abuse of their primary caregiver. About 18% of the youth said their parents were divorced, and about 8% said somebody in their household or family was put in jail. Results were assessed. This outcome has demonstrated that ACE-exposed adolescents would more likely spend time in peer environments known to be associated with increased substance use.

2.24 (Ashna Arora, 2023)

It explores the potential of programs being implemented across the country by police departments in an effort to redirect people possessing narcotics from prosecution to treatment and recovery. This is done by applying the eligibility criteria and the staggered narcotics arrest diversion implementation across Chicago during the period from 2018 to 2020. The difference-in-differences framework—with the staggered implementation of the program across the Chicago districts and individual-level eligibility criteria for the diversion—will be applied in estimating the effect of diversion on eligible individuals. The empirical design will source data from the Chicago Police Department, Thresholds Data, and a survey of police officers. This can be done to serve the purpose of assessing the eligibility for diversion, connections that will subsequently be formed with the substance use counselors, and subsequent arrests post-diversion.

The narcotics arrest drug diversion program in Chicago worked magic in facilitating linkage to substance use treatment and reducing further arrests. This program was capable of making contact with those medically diagnosed as having a substance use disorder and reducing the chances of recidivism, ultimately increasing the cessation rate of substance use.

2.25 (Azmawati Mohammed Nawi, 2021)

The systematic review deals with the uses of drugs among different ethnicities within Chicago City suburbs. The paper will look at drug use patterns in different ethnic groups, such as African Americans, Hispanics, and Whites, residing in the Chicago City suburbs. The methodology used in this paper are Systematic review using three databases: PubMed, EBSCOhost, and Web of Science; Preferred Reporting Items for Systematic Reviews and Meta-Analyses checklist

and Search strategy with verified and validated keywords. Quality assessment was performed using the 2018 edition of the Mixed Methods Appraisal Tools. A number of them are methodology using existing research articles.

The individual risk factors of the study include high impulsivity, rebelliousness, bad emotional control, low religiosity, pain catastrophizing, homework completeness, total screen time, alexithymia, maltreatment or bad upbringing, psychiatric disorders, e-cigarette exposure history, behavioral addiction, perception of low risk, high accessibility perception surrounding drugs, and highly positive attitude towards synthetic drugs. These family risk factors included prenatal maternal smoking, poor maternal psychological control, low parental education, negligence, poor supervision, uncontrolled personal money, and the presence of family members who abuse substances. Some of the community risk factors mentioned included having acquaintances who use drugs. Individual protective factors against drug abuse included optimism, high mindfulness, social anxiety, strong beliefs against substance abuse, desire to maintain one's health, high level of paternal awareness of drug abuse, school connectedness, structured activity, and strong religious beliefs. The study also discovered a positive association between a history of maltreatment and drug abuse among adolescents, particularly among females.

2.26 (Raymond L. Moody Jr, 2022)

Patterns of substance use disorders among Black (cisgender sexual minority) men and Black (transgender women) participants residing in Chicago during the first wave of the COVID-19 pandemic were studied in this paper. The methodologies adopted include chi-square tests and modified poisson regression models.

The results of the study found that more than 70% of the samples reported alcohol consumption at least once a day. Additionally, more than 70% reported marijuana consumption in the last 14 days. Alcohol and Marijuana were found to be the most common substances that were misused. It was also found that about 60% of the samples engaged in polysubstance use. Very minor differences were found in the patterns of substance misuse of the participants but those were due to different socioeconomic stature.

2.27 (Basmattee Boodram, 2022)

This study is conceived and led in enlightening the pattern of drug use among people who inject drugs residing in both the city of Chicago and surrounding suburban areas. The material collected in the research has been applied with the use of the Meta-Analysis application, in the study of demographical features, drug of choice, and network estimates of PWID within the Chicago area. They used Exponential Random Graph Models for modeling the dependencies of behavioral parameters, especially in modeling the syringe sharing networks among the PWID; and Computational models to understand the interaction between the interplay of factors that influence the drug use behaviors and hepatitis C transmission among the PWID in the metropolitan city of Chicago.

The study reported that male PWID made up about 68% of the population in the greater Chicago area. Non-Hispanic white individuals made up 53% of PWID, followed by Hispanic individuals, 23%, and non-Hispanic Black individuals, 14%. Heterosexual injection drug use was common, 85%, compared with gay/homosexual, 3%, and bisexual, 11%. Heroin was also the most commonly injected drug among the PWID, 98% of participants reporting its use. Cocaine injection was reported by 36% of PWID, and "speedball" and methamphetamine use by 24% and 3% of participants, respectively. Network data revealed how syringe sharing occurred between different groups.

2.28 (Rebecca S. Bolinski, 2022)

The paper describes changes in drug availability, use behaviors, and harm reduction services in rural environments. The data from this study describe how external influences, such as the pandemic, shape drug use patterns, and it is important to understand these dynamics in different populations. In this study, a mixed-method approach was utilized, with both in-depth interviews of a qualitative nature carried out to draw critical knowledge regarding COVID-19, community impacts, attainment and use of drugs and a survey, covering questions relating to demographics, drug availability and use patterns, sharing practices, and symptoms of mental health amongst the participants.

The pandemic markedly increased the misuse of fentanyl as access to heroin decreased and the price for methamphetamine went up. Participants reported more willingness to report using alone during the pandemic compared to before the pandemic and also reported more frequently witnessing or experiencing an overdose. Mental health concerns like anxiety and depression had also been found to have increased during the pandemic. Participants said a mobile harm reduction model provided them with some key preventive measures: fentanyl test strips to check drugs. They also claimed that naloxone reversed some drug overdoses.

2.29 (Zhiyue Xia, 2023)

Authors of the study have outlined in this paper insights into the spatiotemporal patterns of opioid-related mortality and the influence of COVID-19 on drug overdose deaths in the urban settings. The method of research in the study is via the Spatiotemporal Random Forest Modeling approach. The spatiotemporal patterns, demographic factors, and pandemic-related variables were used in understanding the impact on the opioid mortality rates. An attempt was made at estimating the impacts of the pandemic on opioid-involved deaths by conducting a counterfactual analysis. Data used for the research have been derived from various datasets.

The study has found a higher rate of opioid-involved deaths during the pandemic. It was noted that counties with higher income inequality had higher drug overdose rates in the year 2020. In the modeling of opioid-involved overdose deaths during COVID-19, demographic variables, like race and ethnicity, were found critical indicators. In the final sense, the number of opioid-involved overdose fatalities had a positive correlation with COVID-19-related factors, such as changes in stay-at-home behaviors and home-dwelling time, over time and space.

2.30 (Peter S. Larson, 2022)

The research article evaluated the temporal patterns in the impact of the COVID-19 pandemic on death rates from mental health- and substance abuse-related causes of death in Michigan. The study was based on an interrupted time series and generalized additive poisson regression models. This article investigated three broad categories of deaths caused by mental health. The categories were suicide, drug overdose, and alcohol-related liver failure. It tried to make sense of the shift in temporal patterns for those different causes of death between before and after the pandemic.

In the urban residents, suicides dropped further sharply after the start of the pandemic compared to the rural residents and remained at a low for the rest of the year. In the analysis by age, an increase in suicide incidence in ages above 65 years, compared to those between 18-65 years, was further observed after the beginning of the pandemic. There was also an observed increase in the incidence of drug overdose more in males than females after the beginning of COVID-19.

2.31 (Leslie D. Williams E. L.-A., 2022)

It uses such data in exploring the reported changes in risk factors of drug toxicity among PWID in Chicago during the pandemic period. More specifically, this paper assesses the change in manifestation for psychological symptoms and access to care, and how these changes in risk-bearing factors affect the experience of economic adversity and their relation to the pandemic. The methodology adopted for the study also included the recording of the interviews with the subjects. Next, the alleged financial trial experiences, measured against overdose risk factors during the pandemic with bivariate cross-tabulations, were compared. Adjusted logistic regression analyses for sociodemographic confounders and other variables have also been factored into the study.

Most participants reported mental health issues related to the pandemic, and around half of the sample used more medications alone. The accommodation situation had an interaction that was markedly associated with overdose-related concerns since the beginning of the pandemic. Accommodation was marginally associated with a change in drug integrity/quality. It connected enhanced overdose danger among individuals who infuse drugs to financial difficulties all through the pandemic, which thus influenced those factors at both the behavioral and condition levels associated with dangers of overdose.

2.32 (Fern R. Hauck, 2022)

This is a valuable study on the effects of maternal substance use on infant health outcomes within a predominantly Black population of Chicago. Data were collected for the Black infants who died from SIDS and matched controls for race, age, and weight. Other statistical analyses were also applied.

Smoking of the mothers during and after pregnancy was found to have been related to a significantly higher risk of SIDS. The risks for SIDS were also linked to the use of alcohol and marijuana by mothers. There were significant positive interactions of bed-sharing with the variables of maternal smoking, both during and after pregnancy: alcohol use, and cocaine use.

2.33 (Mary Ellen Mackesy-Amiti, 2022)

In this research the authors have explored the network characteristics of individuals who inject drugs in the Chicago metro area, focusing on their injection, sexual and social support networks. The methodology used in the study includes ongoing longitudinal network-based research known as the alter-ego study. Other methods of the study include using Likert-type scale to assess injection behavior; adaptive lasso logistic regression, to analyze the associations among the network variables and HCV antibody positive status and injection risk behaviors and computer-assisted interviewer-administered questionnaires and GENSI software for data collection on network characteristics.

HCV antibody frequency differed substantially by race and ethnicity; it was 18% in Hispanic persons, 36% in non-Hispanic white participants, 39% in non-Hispanic black participants and 56% in people of mixed or other races. Also, associations between injection risk behaviors and network characteristics were found. Specifically, a mostly Hispanic support network was associated with reduced odds of receptive syringe sharing and incident HCV infection, and having a Hispanic network with reduced odds of HCV infection.

2.34 (Marjorie C. Gondré-Lewis, 2023)

The current study finds concrete trends in Black and White residents' mortality rates in name states with a high proportion of Black urban residency that have officially declared themselves as hotspots for opioid overdose. Data was derived from several sources, including Offices of the Chief Medical Examiner and Departments of Health of specific states and cities. Statistical analyses of trends were performed with data on the number of deaths related to opioids, heroin, and synthetic opioids over some period. That is, opioid overdose-related deaths were compared with homicides over the same periods of time; and the trend in opioid-related death was determined in large urban areas that have high Black populations. Analyses used the FBI Uniform Crime reports, vital statistics data from the CDC and census data.

Another trend one notices is that the opioid epidemic spread into the communities affected by the urban areas, especially to Black communities. Black Americans were noted to be enduring a staggering toll from opioid overdoses, with disparities in proper treatment, rehabilitation, and health care services. The urban opioid overdose death rates were found to have increased due to the adulteration of drugs with the artificial strong fentanyl. Opioid deaths were noted to have surpassed the homicide cases in some states, including Illinois, Maryland, Michigan, and Washington DC.

2.35 (Maryann Mason, 2023)

This is a qualitative study in which the authors have observed and assessed the perceptions of health-care and harm-reduction service providers about the older adults who use non-medical opioids in Chicago, Illinois. The technique applied to this research to collect the samples by the researchers is snowball sampling, a method of non-random or non-probability sampling. A total of 24 interviews were conducted.

The findings from the study indicate that providers conceptualize the patients as a specific population with distinct behaviors, including those that may protect them from opioid overdose. This population was reported to be difficult patients because symptoms of substance use needed to be sorted out from the symptoms of other illness that they may have had due to their age. Treating this population was also described to be complex. The providers have also mentioned that the patients hide in such behaviors that makes them less visible to outreach.

2.36 (Thomas Coombs, 2023)

In this research, the authors discuss the pervasiveness, tendency, and impact of use of substance among the people who are homeless. Systematic review was the methodology followed by the research. Data were collected through a search of 13 databases, for articles that were dated from 1st January 2007 to 31st December 2020. Search terms used were the same as previous reviews. Data analysis was done using SPSS version 26. Adverse events were categorized into six main systems within the body for comparison. Data reliability was checked by using Higgins's test of homogeneity.

The results revealed that the traditionally popular drug compounds remained popular among people who were homeless. However, NPS had become a favorite amongst this population since 2017. Substance use was found to be higher in males than in females. Both the genders were found to face mental and physical health problems due to the use of drugs. Use of substance among people who were homeless was seen as a cause for many mental health issues. Physical health problems included damage to multiple organs. It was also found that the rise of NPS resulted in a declining use of other popular drugs.

3. Methodology

This research paper methodology section envisages the framework of methods used in conducting a detailed analysis of drug abuse in different ethnicities in the US. Because this research is done from Bangladesh and involved geographical constraints, the paper primarily relies on secondary data. In a way, it enables a broad review of data and the synthesis of existing literature on the topic. On the negative side, the data may not be updated and from the selection and interpretation of the data, it may as well be biased. This strategy, based on relying on established and validated sources, used a systematic literature search approach, elaborated selection criteria, data extraction procedures, and analytical methods to gain meaningful insight on the research topic.

3.1 Strategy for Literature Search

3.1.1 Utilized Databases: A systematic investigation of digital databases such as Google Scholar and Research Gate was conducted to locate relevant literature.

3.1.2 Search Phrases and Keywords: The right combination of search phrases and keywords have been utilized in the search process to maximumly capture the essence of the subject under investigation. Some of the used search phrases included "drug", "drug abuse", "drug use", "substance", "substance misuse", "ethnicity", and "United States", and others, all of which has plenty of variations of the terms under research. Therefore, the selection and combination of this search terms were to ensure that no stone is left unturned in the quest for relevant literature.

3.1.3 Inclusion/Exclusion Criteria: Only studies that passed stringent criteria were included in this study. The criteria require unique research finding of the study in terms of drug addiction or substance misuse amongst ethnic groups in the USA, to be published under peer-reviewed publications, and to be written in English. Reviews or comments were excluded or if they failed to explicitly focus on the USA context. Such a rigorous selection process ensures only the best quality studies are available to evaluate.

3.2 Selection Criteria

3.2.1 Title and abstract screening: The objective of the screening was to identify publications that may be relevant to the topic of the study. Assessment of full texts of potentially relevant publications against pre-specified eligibility criteria determined inclusion.

3.3 Data extraction

3.3.1 Data Collection: The included publications gathered data using a standardized form. The data collected included the ethnic group of the study subjects, the drugs they specifically misused, the prevalence, factors that escalated the risk of substance misuse, and any other findings relevant to substance misuse for specific ethnic groups.

3.3.2 Data Synthesis: The quantitative data from studies that were included in the review were pooled using meta-analytic techniques whenever possible. This was done by computing the effect sizes, statistical testing, and even comparisons of prevalence rates. The effect sizes were computed with an intention to compare prevalence rates of drug misuse between different ethnic groups. This was done by first reviewing the data, the key findings, and calculating the effect sizes using the preset formulas. The subgroups analyzed were to compare the different ethnic groups to ascertain whether there were any differences in the patterns of drug misuse between them. The process of synthesizing the qualitative data, and specifically the thematic findings, used a narrative review approach to identify the emerging themes and patterns. The factors considered in the data were the study design, representativeness of the sample, validity of the measurements, and likely sources of bias. This is followed by a synthesis and closer examination of the data, key findings identification, and developing a coherent narrative in which findings from across the disparate studies are integrated.

3.4 Ethical Considerations

3.4.1 Ethical Compliance: This one is a secondary form of research, based on readily available data that was derived from previous research. As such, no ethical approval was required, given that I had no need to engage in the collection of raw data or respondent questioning, or analyze materials. Academic integrity has also been maintained at all levels of proper citation of original works.

4. Analysis

The use and abuse of substances have presented many challenges across the United States and affect not only individuals but all different ethnicities and communities. The studies of drug abuse among ethnic groups show inequalities, barriers, and complex determinants for the variation in different misuse rates. This paper presents an analysis of the findings from studies conducted on substance misuse among minority communities, showing the higher impact of abuse within foreign communities. An analysis of the interplay of race, socioeconomic status, access to health services, and criminal justice policies can help in bringing out unique dynamics to which the minority communities are subjected to the context of substance misuse. It is very important to learn these dynamics better in order to develop interventions and support systems to help deal with the causes and effects of substance abuse and promote better health equity across ethnic populations of the USA.

4.1 Role of age, gender, and adverse childhood experience (ACE) in Substance Abuse:

- **Adolescents:** According to the research, the adolescent is a vulnerable stage for drug abuse. In some papers, youth have reported using substances, drinking, using illicit drugs, and cigarette smoking.
- **Young Adults:** As reported in previous research, the effect of the urban environment on trajectories of marijuana during initial adulthood means that those residing in or around neighborhoods with higher levels of crime, alcohol shops, and poverty at age 18 are associated with continued marijuana use during emerging adulthood.
- **Adults:** A study focused on the beginning phase of the Covid-19 pandemic in Michigan has reported an increased rate of mortality through suicide, drug overdose, and liver-failure due to alcohol consumption. In addition, in the context of colleges, it is reported that males tend to use substances more than females.
- **Gender Differences in ACEs and Substance Use Patterns:** Finally, adverse childhood experiences were found to be very important in the context of substance abuse. Reports indicated that adverse experiences were more common among girls, particularly psychological abuse and family substance abuse. In contrast, boys were more likely to experience physical abuse.

4.2 Socioeconomic and environmental factors:

Places characterized by socio-economic disadvantage, more competitive labor markets, and easier access to prescription opioids experienced relatively more drug-induced deaths. Social stigma, for the most part, keeps drug users away from treatment options and harm reduction services.

Higher rates of substance abuse, for example, overdose of prescription opioid and heroin, corresponded to an increase in poverty in families. Unemployment was reported to be directly proportional to the increase in overdose of drugs like heroin. The average of household

income was inversely related to the rates of overdose; the higher the income, the lower the rates of POD and HOD. The distribution of deaths due to overdose involving fentanyl in Cook County was unique, where specific areas in Cook County were at higher risk for overdoses involving fentanyl. It is this that provides the substantial association between neighborhood economic hardship and deaths from overdose varied by race/Hispanic origin subgroup.

In some instances, geographic exclusion resulted in higher heroin accessibility for Blacks in urban setups in comparison to white, while concomitantly reducing access to resources such as transport, employment and education.

5. Findings:

5.1 Disproportionate Effect on Minority Communities:

Abusing substances differentially burdens Hispanic and Black communities compared to Non-Hispanic and Whites in the US. Indications are a higher occurrence of substance abuse and associated consequences for minority populations. In general, such inequalities in criminal justice response to research results are based on evidence that people in minority groups have been disproportionately at the receiving. Many of the structural problems left a legacy of substance abuse and incarceration, where the minorities are overrepresented in the criminal justice system. As the data shows, there is a serious and continuous need for structural change in criminal justice policy reform and culturally appropriate access to substance abuse treatment if this current, disproportionate burden of misuse and incarceration among minority communities is to be reduced.

5.2 Issues Faced by Minorities:

The major challenges faced in the minority communities are related to access to rehabilitative treatment. The huge barrier noted is the fear of incarceration that deters people from seeking much-needed support against substance abuse problems. This fear is embedded in historical and ongoing inequalities within the criminal justice system, especially for people of color. This data set reveals that such fear of penal procedures may keep people from minority communities away from seeking necessary help in solving substance misuse.

Moreover, the socioeconomic factors become pivotal in building blockades for minorities seeking help from rehabilitative treatment centers. It is found that inequalities in access to affordable healthcare contribute to the challenges faced by the minority communities in receiving necessary treatment services. Barriers to healthcare resources, such as mental health and substance abuse treatment programs, may delay or prevent people from getting timely and appropriate treatment for their substance abuse issues. The inequalities faced in healthcare further add up to the preexisting challenges faced by the people of color in the USA.

Overall, the data studied shows a clear picture of the interplay of the obstacles faced by minority groups in healthcare, socioeconomic environment, and jurisdiction. Overcoming the aforementioned barriers will require a comprehensive strategy that serves not only to fear of incarceration but also to socioeconomic differences constraining access to necessary healthcare facilities for everyone and ensuring equal and proper treatment access.

5.3 Association between Victimization and Substance Use:

The most important finding is the correspondence of the experience of victimization and substance-using behaviors among people (sexual and gender minority) assigned female at birth. The findings suggest that substance-using behaviors for this population are most likely to be found in those who have been victimized. This connotation reveals the impact of traumatic experiences, such as victimization, on the patterns of substance use within specific groups.

Furthermore, the findings point out the link between victimization and behaviors of substance use as an important illustration of the interaction between social factors, mental health, and substance misuse in this population group. It makes imperative the call for targeted interventions and support services that can address both the underlying trauma of victimization and the substance use issues faced by individuals (sexual and gender minority) assigned female at birth.

The data studied in this research collectively provides evidence that victimization experiences may play a huge role in substance misuse among this population. Diagnosing the effect of victimization on substance use within this certain population can be a useful method to guide tailored interventions that propose holistic support for the trauma of victimization and substance misuse issues, having better outcomes for the people in this demographic group.

5.4 Ethnicity in Substance Abuse:

In this research, discrimination, and substance abuse have been found to be correlated in many cases. The data studied shows that discriminative experiences, whether based on race, ethnicity, or other aspect, might increase the vulnerability of minority ethnic groups towards substance abuse. Discrimination, indeed, brings stress, trauma, and relegation to people and, hence, might drive an individual to cope with those negative experiences with the use of substances.

Additionally, such a relationship between discrimination and substance abuse denotes the broader social factors of health that affect the patterns of substance use among ethnic communities. The prevalence of substance abuse in different ethnic communities can be dealt with by forming appropriate strategies that prevent and reduce substance misuse among populations only by facing discrimination and advancing social equity. Understanding the relationship between discrimination and substance abuse among different ethnicities can enable the tailoring of interventions to target underlying social factors that might result in substance use. Providing support, cultivating cultural sensitivity, and fighting discrimination are part of the steps that should be taken in mitigating social inequalities on substance abuse within diverse

ethnic communities. Understanding the call, such kinds of information emphasize the need to have a holistic approach that considers the intersection of social determinants, such as discrimination, in the explanation of the substance abuse disparity between diverse ethnic backgrounds.

5.5 The effect of Opioid Epidemic on Black and Hispanic Communities:

It is noted that Hispanics, who constitute about 17% of the U.S. population, amount to 50% of all Federal Drug cases (Jasmine Drake, 2020). This means that the share of drug-related legal cases that affect the Hispanic communities is disproportionately huge. As such, being that the differential between the two percentages of drug traffickers is so much larger than that of the general American population, one can ultimately reach the understanding that the drug laws and the means by which the law enforcement agencies process drug cases and alleged drug offenders are structurally biased. In particular, such a gap could be due to the compound effect of biased law enforcement practices, differences in legal representation, and unequal treatment by the court system.

The opioid crisis has been so devastating, resulting in deaths related to opioids among Blacks and Hispanics. Opioid-related deaths in Blacks and Hispanics are disproportionately high when compared to non-Hispanic Whites. Such gaps point to differences in healthcare access, treatment availability for addiction, and prevention. Some of these disparities could be because of socio-economic factors, lack of access to quality health care, stigma associated with drug abuse, and history of disparities in health care provision. The data found underscores the needs for targeted efforts and flexible laws that, in particular, take into account the worries of Black and Hispanic populations affected by the opioid crisis.

Because opioid-related harm and overdose deaths are problems with disparities that need to be eliminated, this places a special focus on the role that reducing the disparities experienced by racial and ethnic minorities has in the realization of this objective, including decreased access to treatment, culture-driven help-seeking habits, and engagement with the socioeconomic determinants of health. In recognizing these important issues and actively remedying the disparities currently in place, providers, policymakers, and community organizations can work to design and implement equitable, effective opioid epidemic interventions in support of the health and well-being of Black and Hispanic individuals, who are disproportionately affected by issues of substance abuse.

5.6 Neighborhood and Economic Factors:

Furthermore, this study also points out that neighborhood and economic variables are very strong mediators of the impacts due to drug misuse, in particular mortality from overdose due to opioids. This study reveals elevated rates of opioid-related deaths in resource-starved, underinvested areas, with distinct disparities among different ethnic and Hispanic origin subgroups. This shows how neighborhood conditions and the outcomes of substance misuse

are correlated and further reveals how inequalities in investment and community resources could raise the risk for opioid-related harm, mainly in underprivileged groups.

The statistics also bring to light the relationship between the overdose of opioids and the problems of neighborhood economic adversities. That points out how these factors in the economic sectors influence the trends of substance use disorders in subgroups of different racial and Hispanic origins. Economic adversities, such as unemployment, poverty, or limited access to essential services, are conditions which tend to aggravate problems of substance use disorder and increase vulnerability to harm from opioids.

Studying elaborately about how economic and neighborhood conditions affect drug misuse outcomes can acknowledge the root causes of these disparities in opioid-related harms, in order to design treatments that target the underlying socioeconomic determinants. There is a lot that can be done to reduce substance abuse, prevent it from occurring, and reduce opioid-related overdose deaths with measures to strengthen community resources, promote economic stability, and improve access to care and addiction treatment services.

6. Limitations

6.1 Publication Bias: Though I have tried my level best to retrieve all the published literature, I am aware that the results may be influenced by publication bias and the limitations of the critical studies from which it is taken for review.

6.2 Geographical Limitations: There are few geographical limitations that come in the way of my research, for instance, some of the databases are not available to me, real-time sources of information are not available, and so on. It might result in either a lack of current information or in some biased representation of data. Furthermore, my reliance on the published literature has restrained me from exploring emerging trends or capturing the firsthand observations from the study populations. These are points to bear in mind while interpreting the results of this study, as they may affect the generalization and applicability of the findings.

7. Recommendations:

7.1 Unbiased Access to Treatment and Support Services:

It is found in the research that the accessing support or treatment is blocked by the lack of affordable healthcare and fear of incarceration in ethnic communities. Therefore, it is inevitably significant to remove these barriers to ensure equal health outcomes and address substance abuse issues among the people of minority races.

Interventions need to be culturally sensitive and community-based in order to be effective with the needs and interests of these diverse ethnic populations. By recognizing the cultural nuances and challenges faced by people of color, it is possible to create specific support that is sensitive to cultural, social and economic contexts. All in all, culturally competent care can build trust and reduce stigma while improving the engagement of minority individuals in substance abuse treatment and support services.

Community interventions may include outreach, peer support programs, education, culturally appropriate material, and last but not least, partnerships with other local organizations to ensure access to their services and help them make the needed changes so that the services would be appropriate for the needs of the minorities. Hence, healthcare providers and policymakers can come together to reduce these inequalities in access to treatment and support services by including cultural sensitivity and community engagement in their priorities to achieve better health outcomes and well-being among the minority populations most affected by the substance abuse issues.

7.2 Policy Reforms and Criminal Justice:

Ethnic communities are at greater risk of being arrested for drug-related offenses and being incarcerated for longer periods than Whites. Thus, policies that cease such current inequitable practices and institute fair sentencing guidelines can lead to the reduction of the lopsided burden that ethnic minority populations currently bear in the criminal justice system.

A better way of getting rid of such inequalities may be to rehabilitate and support rather than punish. Diversion from prison to treatment programs, counseling, and other community services might just be a means to try to get to the bottom causes of substance abuse and offer clients resources to overcome addiction. Treatment in this approach is therefore based on recovery and the well-being of offenders; the approach, by so doing, contributes to a reduction in recidivism rates while providing healthy communities.

Overall, these reforms and alternatives to incarceration suggest that the fair and effective way to deal with substance abuse is to advocate for all people receiving support and resources to be able to cope with addiction and towards healthier living.

7.3 Prevention and Education Programs:

Targeted prevention and educational programs regarding substance use in minority communities will be designed in an effort to reduce substance abuse problems in ethnic populations in the USA. The developed programs may target at the social determinants of health and systemic issues unique to these populations. Thus, prevention and education efforts would be more attuned to the particular needs and situations that surround minority communities to accomplish efficacy in interventions that make sense and address the very roots of substance abuse.

Also, raising awareness that focuses on the hub of such issues as the epidemic of opioid among Black and Hispanic communities is crucial to garner resources and support for the cause of prevention. Such identification of differences may help stakeholders advocate for targeted interventions, funding, and policy changes to address these disparities. Such awareness of the specific challenges and vulnerabilities garners support from the implementation of proof-based inhibition strategies and programs for treatment by policymakers, healthcare providers, and community organizations.

7.4 Research and Data Collection:

Additional studies are required for offering more impactful interventions that are sensitive to all cultures and meet the needs of every racial group. Efforts on data collection that emphasize the intersection of race, ethnicity, socioeconomic status, and others influencing substance abuse need to be prioritized in diverse populations.

By looking at the interaction of the social determinants of health, researchers could establish patterns and disparities in substance abuse among various ethnicities more clearly. This approach would help to bring into light deeper drivers, such as economic hardship, access to resources, and systemic inequalities, and interventions could take a comprehensive approach to treating such multifaceted issues.

The approach will inform health equity and social justice through evidence-based policies and programs, addressing the underlying determinants of substance abuse disparities among ethnic communities.

Ongoing research, greater public awareness, and building resources to expand the scope of prevention will therefore be important to improve the scale of prevention and increase the improvement in outcomes for people with substance abuse problems in minority communities. And so, we will be able to have a more just and efficient approach to substances with health and resilience for all people, regardless of their origin and ethnic background.

8. Discussion

The key and vital aim of the study was to unearth prevalence of substance abuse within the complex setting of ethnic diversity in the US. This study was particularly focused on unveiling the subtle differences in drug use behaviors between these communities, motivated by the prevalence of drug abuse across several ethnicities. Central in this paper were, therefore, some critical questions that needed to be understood in respect to the varied causes of substance misuse among these varied ethnicities. These causes cut across a wide variety of influences, among them being social economic status, cultural norms, historical policies, and access to health services. The influence of such factors had been found in previous studies on the topic as well. Communities that live on low-income and that have lesser access to basic needs are severely and disproportionate affected by the alarming substance misuse issues (Veronica A. Peara, 2018). Financial challenges and many disruptions caused by the pandemic in 2019 has massively increased the consumption of substances, mostly among people to inject substances. (Leslie D. Williams E. L.-A., 2022).

Furthermore, aligned with former researchers, this study focused on identifying and explicating the barriers to accessing treatment for substance use problems in minorities, from systemic inequalities within the healthcare system, to fear of legal ramifications and historical injustices that have limited treatment availability and utilization by people from diverse racial backgrounds. Enrollment in employment, medical coverage, and financial support reentry programs was more common for respondents living in a more cohesive community. Whereas those who returned to disordered neighborhoods-defined all too often by poverty and residential mobility-were not only bereft of networks, they were also less likely to enroll in reentry programs. (Lin Liu, 2021).

The paper sought, in addition, to assess the deep effect of social elements on patterns of drug abuse in diverse ethnic communities. These, in turn, emanate from a wide range of social factors, including discrimination, economic adversities, neighborhood conditions, and access to resources, which have an influential effect on the definition of substance use behaviors and outcomes among members of varied ethnic backgrounds.

The study, therefore, aimed to bring better insight into substance misuse trends and disparities, informing the development of targeted interventions and policies to ensure health equity and resilience across different ethnic populations.

9. Conclusion

In conclusion, complex drug use behaviors and disparities exist within varied communities. The findings of this study highlighted the disproportionate burden of substance abuse on the Black and Hispanic residents compared to White and non-Hispanic. Indicative is the all-pervasive impact of historical policies and systemic inequalities in the drug misuse patterns of the population. There are very strong barriers to rehabilitative treatment, from fear of incarceration to healthcare disparities, that prevent essential support for substance abuse issues in minority populations. Furthermore, the study places immense significance on the critical role that social determinants play in shaping substance misuse disparities across varied ethnic backgrounds. Advocating for structural changes in criminal justice policies, advocating for economic stability, and enhancing access to care and addiction treatment services will help minimize these disparities in ethnic communities when it comes to substance abuse outcomes. Only through raising public awareness, expanding prevention efforts, and building resources that help people struggling with substance abuse within minority communities can a just and fair approach to substance abuse problems be established while promoting the health and resiliency of all people, regardless of their ethnic background.

This approach, developed in line with the insights and recommendations gleaned from the current analysis, might spur meaningful changes and improvements in the outcomes that people from different ethnic backgrounds experience in the U.S. while dealing with their substance abuse problems. It is only in the increased awareness of the public, broadened efforts at prevention, and resources that assist individuals battling substance misuse in minority communities that an approach to the misuse of substances that is more just and fair can be established, and health and resiliency achieved for all individuals, no matter their ethnic background.

Although a vast area of the topic has been covered, further research is needed for understanding the broader landscape of substance misuse in the USA and mitigating this major health concern.

References

- Anna Hotton, M.-E. M.-A. (2021). Trends in homelessness and injection practices among young urban and suburban people who inject drugs: 1997-2017 . *Drug and Alcohol Dependence*.
- Ashna Arora, P. B. (2023). Policing Substance Use: Chicago's Treatment Program for Narcotics Arrests . *Chicago's Treatment Program for Narcotics Arrests*.
- Azmawati Mohammed Nawir, R. I. (2021). Risk and protective factors of drug abuse among adolescents: a systematic review. *BMC Public Health*.
- Basmattee Boodram, M. E.-A. (2022). People who inject drugs in metropolitan Chicago: A meta-analysis of data from 1997-2017 to inform interventions and computational modeling toward hepatitis C micro elimination. *PLOS One*.
- Beth A. Reboussin, N. S.-H. (2018). The Impact of the Urban Neighborhood Environment on Marijuana Trajectories During Emerging Adulthood. *Prevention Science*, 20(10.1007/s11121-018-0915-4), 270–279.
- Beth A. Reboussin, N. S.-H. (2018). The Impact of the Urban Neighborhood Environment on Marijuana Trajectories During Emerging Adulthood. *Prevention Science*, 270–279.
- Cara Jane Bergo, J. R. (2021). A Vulnerability Assessment for a Future HIV Outbreak Associated with Injection Drug Use in Illinois, 2017–2018 . *Front. Sociol.*
- Elizabeth D. Nesoff, P. C. (2019). The Geographic Distribution of Fentanyl-Involved Overdose Deaths in Cook County, Illinois. *RESEARCH AND PRACTICE*.
- Fern R. Hauck, S. R. (2022). Maternal Smoking, Alcohol and Recreational Drug Use and the Risk of SIDS Among a US Urban Black Population . *Frontiers in Pediatrics*.
- Gregory Phillips II, D. F. (2020). Engagement with LGBTQ community moderates the association between victimization and substance use among a cohort of sexual and gender minority individuals assigned female at birth. *Addictive Behaviors*.
- Hanlin Zhou, L. L. (2021). Using Google Street View imagery to capture micro-built environment characteristics in drug places, compared with street robbery . *Computers, Environment and Urban Systems* .
- Harvey L. Nicholson, J. V. (2018). Gender Differences in Prescription Opioid Misuse Among U.S. Black Adults. *Substance Use & Misuse*, 639-650.
- Jacqueline Korpicsa, A. S. (2020). 2.13 Declining health risk exposure among Chicago public high school students: Trends from the Youth Risk Behavior Survey 1997–2017 . *Preventive Medicine Reports*.
- Jasmine Drake, C. C. (2020). Exploring the impact of the opioid epidemic in Black and Hispanic communities in the United States. *Drug Science, Policy and Law*.
- Jeffery Sauer a, K. S. (2021). *Journal of Substance Abuse Treatment*.
- Jerel M. Ezell, P. M. (2020). Stigmatize the use, not the user? Attitudes on opioid use, drug injection, treatment, and overdose prevention in rural communities. *Social Science & Medicine*.
- Karen A. D'Angelo, W. H. (2019). The drug issue really isn't the main problem"— A photovoice study on community perceptions of place, health, and substance abuse. *Health & Place*, 257-264.

- Leslie D. Williams, E. L.-A. (2022). Economic Challenges and Behavioral and Mental Health Risks for Overdose during the COVID-19 Pandemic among People Who Inject Drugs. *International Journal of Environmental Research and Public Health*, 19.
- Leslie D. Williams, M. E.-A. (2021). Drug use-related stigma, safer injection norms, and hepatitis C infection among a network-based sample of young people who inject drugs. *Drug and Alcohol Dependence*.
- Leslie D. Williams, P. U. (2020). Trends over time in HIV prevalence among people who inject drugs in 89 large US metropolitan statistical areas, 1992-2013. *Annals of Epidemiology*, 12-23.
- Lin Liu, C. A. (2021). The Roles of Family, Community, and Services in the Prevention of Illicit Drug Use: Findings From a Sample of Released Prisoners. *Journal of Drug Issues*, 354–371.
- Lindsay Lezana, C. L. (2020). The role of gender in the relationship between adverse childhood experiences and delinquency and substance use in adolescence. *Journal of Criminal Justice*.
- Marika Sigal, B. J. (2021). Neighborhood, Peer, and Parental Influences on Minor and Major Substance Use of Latino and Black Adolescents. *Children*.
- Marjorie C. Gondré-Lewis, T. A.-L. (2023). The Opioid Epidemic: a Crisis Disproportionately Impacting Black Americans and Urban Communities. *Journal of Racial and Ethnic Health Disparities*, 2039–2053.
- Martin, C. C. (2019). High Socioeconomic Status Predicts Substance Use and Alcohol Consumption in U.S. Undergraduates. *Substance Use & Misuse*, 1035-1043.
- Mary Ellen Mackesy-Amiti, J. F. (2022). Egocentric network characteristics of people who inject drugs in the Chicago metro area and associations with hepatitis C virus and injection risk behavior. *Harm Reduction Journal*.
- Maryann Mason, L. A. (2023). Health care and harm reduction provider perspectives on treating older adults who use nonmedical opioids: A Chicago case. *BMC Health Services Research*.
- Monnat, S. M. (2018). The contributions of socioeconomic and opioid supply factors to U.S. drug mortality rates: Urban-rural and within-rural differences. *Journal of Rural Studies*, 319-335.
- Nisha Behariera, J. D. (2018). Associations of Adolescent Exposure to Severe Violence with Substance Use From Adolescence into Adulthood: Direct Versus Indirect Exposures. *SUBSTANCE USE & MISUSE*, 191-202.
- Peter S. Larson, R. S. (2022). Impact of the COVID-19 pandemic on temporal patterns of mental health and substance abuse related mortality in Michigan: An interrupted time series analysis. *The Lancet Regional Health- Americas*.
- Raymond L. Moody Jr, Y.-T. C. (2022). Polysubstance use in a community sample of Black cisgender sexual minority men and transgender women in Chicago during initial COVID-19 pandemic peak. *Substance Abuse Treatment, Prevention, and Policy*.
- Rebecca S. Bolinski, S. W.-A. (2022). The Impact of the COVID-19 Pandemic on Drug Use Behaviors, Fentanyl Exposure, and Harm Reduction Service Support among People Who Use Drugs in Rural Settings. *International Journal of Environmental Research and Public Health*.

- Stritzel, H. (2021). Peer and Community Influences on Adolescent Substance Use in the Context of Adverse Childhood Experiences. *Sociological Perspectives*, 1-20.
- Tamara Rushovich, M. A.-A. (2020). Opioid-related overdose deaths by race and neighborhood economic hardship in Chicago. *Journal of Ethnicity in Substance Abuse*, 22-35.
- Thomas Coombs, A. A.-J. (2023). Understanding drug use patterns among the homeless population: A systematic review of quantitative studies . *Emerging Trends in Drugs, Addictions, and Health*.
- Veronica A. Peara, W. R.-A. (2018). Urban-rural variation in the socioeconomic determinants of opioid overdose. *Drug and Alcohol Dependence*, 66-73.
- Zhiyue Xia, K. S. (2023). A counterfactual analysis of opioid-involved deaths during the COVID-19 pandemic using a spatiotemporal random forest modeling approach. *Health and Place*.